

Entanglement between two qubits one of which interacts with a thermal field

Eugene Bashkirov, Michail Mastuygin*

Department of General and Theoretical Physics, Samara State University, 1 Academician Pavlov str., Samara 443011, Russia

Abstract

In this paper, we have investigated the entanglement between two dipole coupled two-level atoms. The model, in which only one atom is trapped in an lossless cavity and interacts with single-mode thermal field, and the other one can be spatially moved freely outside the cavity has been carried out. We have considered the effect of the atomic coherence on the entanglement behavior. We have shown that a thermal field might cause high entanglement between the atoms both for coherent and incoherent initial atomic states only for small values of the cavity mean photon number. In the considered model the atoms would get entangled even when both atoms are initially in the excited state. We have also derived that the degree of entanglement is weakly dependent on the strength of dipole-dipole interaction for coherent initial states.

Keywords: Entanglement, Cavity QED, Negativity, Thermal field, Atomic coherence

1. Introduction

Entanglement between separate quantum systems is one of the key problem in quantum mechanics. It plays a central role in quantum information, quantum computation and communication, and quantum cryptography [1]. Several methods of creating entanglement have been proposed involving trapped

*Corresponding author

Email address: `bash@samsu.ru` (Eugene Bashkirov, Michail Mastuygin)

and cooled ions or neutral atoms in cavities, superconducting circuits, spins in solids etc. [2]. In order to function optimally various applications require maximally entangled states. Because of decoherence, which is generally related to noise, there is great difficulty in generating and keeping the integrity of a pure entangled states. Although the interaction between the environment and quantum systems can lead to decoherence, it may also be associated with the formation of non-classical effects such as entanglement [3]. Thus, understanding and investigating entanglement of mixed states becomes one of the actual problem of quantum information. Recently, Bose et al. [4] have shown that entanglement can always arise in the interaction of an arbitrary large system in any mixed state with a single qubit in a pure state, and illustrated this using the Jaynes-Cummings interaction of a two-level atom in a pure state with a field in a thermal state at an arbitrary high temperature. Kim et al. [5] have investigated the atom-atom entanglement in the system of two identical two-level atoms with one-photon transition induced by a single-mode thermal field. They showed that a chaotic field with minimal information can entangled atoms which were prepared initially in a separable state. Zhang directly generalizes Kim's study to the case when the atoms are slightly detuned from the thermal field, and study how the detuning would affect atom-atom entanglement [6]. Zhou et al. [8] have considered the same problem for nonidentical atoms with different couplings. The entanglement between two identical two-level atoms through nonlinear two-photon interaction with one-mode thermal field has been studied by Zhou et al. [7]. They showed that atom-atom entanglement induced by nonlinear interaction is larger than that induced by linear interaction. In [9] has discovered that two atoms can be entangled also through nonlinear nondegenerate two-photon interaction with two-mode thermal field. The influence of dipole-dipole interaction on entanglement between two cubits induced by one-mode and two-mode thermal field has been investigated in [10]-[13].

The problem of creating or controlling the atomic entanglement is greatly related to the atomic coherence of population between different levels. The authors of papers [14]-[18] have shown that the entanglement between two atoms

induced by one-mode or two-mode thermal field can be manipulated by changing the initial parameters of the atoms, such as the superposition coefficients and the relative phases of the initial atomic coherent state and the mean photon number of the cavity field. They have also discovered that entanglement may be greatly enhanced due to dipole-dipole interaction in the presence of the atomic coherence [17].

While, as has been mentioned in [19] the practical applications in quantum information processing require engineering entangled atoms so this expects operable atoms which can be moved to distance without losing of information. Recently many schemes have been proposed to realize the engineering entangled atoms [20]-[25]. Guo and coauthors have proposed a simple scheme to realize an easily engineered two-atom entangled state. The advantage of this scheme is only one atom is trapped in a cavity, and the other one can be spatially moved freely outside the cavity. But the authors have investigated the atom-atom entanglement only for vacuum initial cavity field. In this paper we study the entanglement dynamics for model supposed in [19] for thermal cavity field taking into account the initial atomic coherence.

2. The model

We consider two identical two-level atoms and one-mode quantum electromagnetic cavity field. The first atom is trapped in a lossless microcavity and resonantly interacts with the cavity field of the frequency ω . The second atom lies beside the first atom out of the cavity. We assume that the distance between atoms can compare with a wavelength on working transition. In this case the dipole-dipole interaction should be included. The Hamiltonian of this system can be written as

$$H = (1/2)\hbar\omega(\sigma_1^z + \sigma_2^z) + \hbar\omega a^\dagger a + \hbar g(\sigma_1^+ a + a^\dagger \sigma_1^-) + \hbar J(\sigma_1^+ \sigma_2^- + \sigma_1^- \sigma_2^+), \quad (1)$$

where $(1/2)\sigma_i^z$ is the inversion operator for the i th atom ($i = 1, 2$), $\sigma_i^+ = |+\rangle_{ii}\langle -|$, and $\sigma_i^- = |-\rangle_{ii}\langle +|$ are the transition operators between the excited

$|+\rangle_i$ and the ground $|-\rangle_i$ states in the i th atom, a^+ and a are the creation and the annihilation operators of photons of the cavity mode, g is the coupling constant between atom and the cavity field and J is the coupling constant of the dipole interaction between the atoms and $|+\rangle$ and $|-\rangle$ are the excited and the ground states of a single two-level atom. The two-atom wave function can be expressed as a combination of state vectors of the form $|v_1, v_2\rangle = |v_1\rangle|v_2\rangle$, where $v_1, v_2 = +, -$.

We consider that the initial states of atoms are the coherent superposition of the two levels, that is,

$$\begin{aligned} |\Psi_1(0)\rangle &= \cos\theta_1|+\rangle_1 + e^{i\varphi_1}\sin\theta_1|-\rangle_1, \\ |\Psi_2(0)\rangle &= \cos\theta_2|+\rangle_2 + e^{i\varphi_2}\sin\theta_2|-\rangle_2. \end{aligned} \quad (2)$$

Here θ_1 and θ_2 denote the amplitudes of the polarized atoms, and φ_1 and φ_2 are relative phases of two atoms, respectively. So the initial density matrix for two-atom system can be written as

$$\rho_A(0) = |\Psi_1(0)\rangle|\Psi_2(0)\rangle\langle\Psi_1(0)|\langle\Psi_2(0)|.$$

The initial cavity mode state are assumed to be the thermal one-mode state

$$\rho_F(0) = \sum_n p_n |n\rangle\langle n|.$$

The weight functions are

$$p_n = \frac{\bar{n}^n}{(1 + \bar{n})^{n+1}},$$

where $\bar{n}_i$ is the mean photon number in the i th cavity mode,

$$\bar{n}_i = (\exp[\hbar\omega_i/k_B T] - 1)^{-1},$$

k_B is the Boltzmann constant and T is the equilibrium cavity temperature.

The initial density matrix of the whole system is

$$\rho(0) = \rho_A(0)\rho_F(0) = \sum_n p_n |\Psi_1(0)\rangle|\Psi_2(0)\rangle\langle\Psi_1(0)|\langle\Psi_2(0)||n\rangle\langle n|, \quad (3)$$

Before considering the dynamics of the system for thermal initial cavity field, it is straightforward to first study the case when the trapped two-level atom interacts with Fock state. Suppose that the excitation number of the atom-field system is n ($n \geq 0$) the evolution of the system is confined in the subspace $|-, -, n+2\rangle, |+, -, n+1\rangle, |-, +, n+1\rangle, |+, +, n\rangle$. On this basis, the eigenfunctions of the Hamiltonian (1) can be written as [19]

$$\begin{aligned} |\Phi_{in}\rangle = & \xi_{in}(X_{i1n}|-, -, n+2\rangle + X_{i2n}|+, -, n+1\rangle + \\ & + X_{i3n}|-, +, n+1\rangle + X_{i4n}|+, +, n\rangle) \quad (i = 1, 2, 3, 4), \end{aligned}$$

where

$$\xi_{in} = 1/\sqrt{|X_{i1n}|^2 + |X_{i2n}|^2 + |X_{i3n}|^2 + |X_{i4n}|^2}$$

and

$$\begin{aligned} X_{i1n} = 1, \quad X_{i2n} = (-1)^{i+1} \frac{S_n}{\sqrt{n+2}}, \\ X_{i3n} = \frac{S_n^2 - (n+2)}{(\sqrt{n+2})\alpha}, \quad X_{i4n} = (-1)^{i+1} \frac{S_n(S_n^2 - (n+2) - \alpha^2)}{(\sqrt{n+1}\sqrt{n+2})\alpha}. \end{aligned}$$

Here $S_n = A_n$ for odd i and $S_n = B_n$ for even i . The corresponding eigenvalue are

$$\begin{aligned} E_{1n}/\hbar = (n+1)\omega + A_n, \quad E_{2n}/\hbar = (n+1)\omega - A_n, \\ E_{3n}/\hbar = (n+1)\omega + B_n, \quad E_{4n}/\hbar = (n+1)\omega - B_n. \end{aligned}$$

Here

$$A_n = \sqrt{W_n + V_n}/2, \quad B_n = \sqrt{W_n - V_n}/2$$

and

$$W_n = 4n + 6 + 2\alpha^2, \quad V_n = 2\sqrt{4(n+1)\alpha^2 + (\alpha^2 + 1)},$$

where $\alpha = J/g$.

To derive the full dynamics of our model one can consider also the basis states $|-, -, 1\rangle, |+, -, 0\rangle, |-, +, 0\rangle$. In this basis the eigenfunctions and eigenvalues of the Hamiltonian (1) are

$$|\varphi_1\rangle = (\alpha^2/\Omega)[|-, -, 1\rangle - (1/\alpha)|-, +, 0\rangle], \quad E_1 = 0;$$

$$\begin{aligned}
|\varphi_2\rangle &= (1/\sqrt{2})[(1/\Omega)|-, -, 1\rangle + |+, -, 0\rangle + (\alpha/\Omega)|-, +, 0\rangle], & E_2/\hbar &= \Omega; \\
|\varphi_3\rangle &= (1/\sqrt{2})[-(1/\Omega)|-, -, 1\rangle + |+, -, 0\rangle - (\alpha/\Omega)|-, +, 0\rangle], & E_2/\hbar &= -\Omega,
\end{aligned}$$

where $\Omega = \sqrt{1 + \alpha^2}$.

At last, the Hamiltonian (1) has one more eigenfunction

$$\varphi_0 = |-, -, 0\rangle$$

which corresponds to energy $E_0 = -\hbar\omega$.

Assume that the whole system is initially in the state $|+, +, n\rangle$ ($n \geq 0$), then at time t , the whole system will evolve to

$$|\Psi(t)\rangle = Z_{11,n}|-, -, n+2\rangle + Z_{21,n}|+, -, n+1\rangle + Z_{31,n}|-, +, n+1\rangle + Z_{41,n}|+, +, n\rangle. \quad (4)$$

Here

$$\begin{aligned}
Z_{11,n} &= e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{41n} X_{11n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{42n} X_{21n} + \\
&\quad + e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{43n} X_{31n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{44n} X_{41n}, \\
Z_{21,n} &= e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{41n} X_{12n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{42n} X_{22n} + \\
&\quad + e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{43n} X_{32n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{44n} X_{42n}, \\
Z_{31,n} &= e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{41n} X_{13n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{42n} X_{23n} + \\
&\quad + e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{43n} X_{33n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{44n} X_{43n}, \\
Z_{41,n} &= e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{41n} X_{14n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{42n} X_{24n} + \\
&\quad + e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{43n} X_{34n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{44n} X_{44n},
\end{aligned}$$

where $Y_{ijn} = \xi_{jn} X_{jin}^*$.

Similarly, when the whole system is initially in the state $|+, -, n+1\rangle$ ($n \geq 0$), then at time t , the whole system will evolve to

$$|\Psi(t)\rangle = Z_{12,n}|-, -, n+2\rangle + Z_{22,n}|+, -, n+1\rangle + Z_{32,n}|-, +, n+1\rangle + Z_{42,n}|+, +, n\rangle. \quad (5)$$

Here

$$Z_{12,n} = e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{21n} X_{11n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{22n} X_{21n} +$$

$$\begin{aligned}
& +e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{23n} X_{31n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{24n} X_{41n}, \\
Z_{22,n} = & e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{21n} X_{12n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{22n} X_{22n} + \\
& +e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{23n} X_{32n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{24n} X_{42n}, \\
Z_{32,n} = & e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{21n} X_{13n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{22n} X_{23n} + \\
& +e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{23n} X_{33n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{24n} X_{43n}, \\
Z_{42,n} = & e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{21n} X_{14n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{22n} X_{24n} + \\
& +e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{23n} X_{34n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{24n} X_{44n}.
\end{aligned}$$

If the initial state of considered system is $|+, -, 0\rangle$, the time dependent wave function takes the form

$$|\Psi(t)\rangle = Z_{12}|-, -, 1\rangle + Z_{22}|+, -, 0\rangle + Z_{32}|-, +, 0\rangle, \quad (6)$$

where

$$Z_{12} = -\frac{i}{\Omega} \sin \Omega t, \quad Z_{22} = \cos \Omega t, \quad Z_{32} = -\frac{i\alpha}{\Omega} \sin \Omega t.$$

For initial state $|-, +, n+1\rangle$ ($n \geq 0$) the time-dependent wave function is

$$|\Psi(t)\rangle = Z_{13,n}|-, -, n+2\rangle + Z_{23,n}|+, -, n+1\rangle + Z_{33,n}|-, +, n+1\rangle + Z_{43,n}|+, +, n\rangle. \quad (7)$$

Here

$$\begin{aligned}
Z_{13,n} = & e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{31n} X_{11n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{32n} X_{21n} + \\
& +e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{33n} X_{31n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{34n} X_{41n}, \\
Z_{23,n} = & e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{31n} X_{12n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{32n} X_{22n} + \\
& +e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{33n} X_{32n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{34n} X_{42n}, \\
Z_{32,n} = & e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{31n} X_{13n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{32n} X_{23n} + \\
& +e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{33n} X_{33n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{34n} X_{43n}, \\
Z_{42,n} = & e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{31n} X_{14n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{32n} X_{24n} + \\
& +e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{33n} X_{34n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{34n} X_{44n}.
\end{aligned}$$

If the initial state of considered system is $|-, +, 0\rangle$, the time dependent wave function of the whole system takes the form

$$|\Psi(t)\rangle = Z_{13}|-, -, 1\rangle + Z_{23}|+, -, 0\rangle + Z_{33}|-, +, 0\rangle, \quad (8)$$

where

$$Z_{13} = -\frac{\alpha}{\Omega^2}(\cos \Omega t - 1), \quad Z_{23} = -\frac{i\alpha}{\Omega} \sin \Omega t, \quad Z_{33} = -\frac{1}{\Omega^2}(1 + \alpha^2 \cos \Omega t).$$

For initial state $|-, -, n+2\rangle$ ($n \geq 0$) we have at time t that

$$|\Psi(t)\rangle = Z_{14,n}|-, -, n+2\rangle + Z_{24,n}|+, -, n+1\rangle + Z_{34,n}|-, +, n+1\rangle + Z_{44,n}|+, +, n\rangle. \quad (9)$$

Here

$$\begin{aligned} Z_{14,n} &= e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{41n} X_{11n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{42n} X_{21n} + \\ &\quad + e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{43n} X_{31n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{44n} X_{41n}, \\ Z_{24,n} &= e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{41n} X_{12n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{42n} X_{22n} + \\ &\quad + e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{43n} X_{32n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{44n} X_{42n}, \\ Z_{42,n} &= e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{41n} X_{13n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{42n} X_{23n} + \\ &\quad + e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{43n} X_{33n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{44n} X_{43n}, \\ Z_{44,n} &= e^{-iE_{1n}t/\hbar} \xi_{1n} Y_{41n} X_{14n} + e^{-iE_{2n}t/\hbar} \xi_{2n} Y_{42n} X_{24n} + \\ &\quad + e^{-iE_{3n}t/\hbar} \xi_{3n} Y_{43n} X_{34n} + e^{-iE_{4n}t/\hbar} \xi_{4n} Y_{44n} X_{44n}. \end{aligned}$$

If the initial state of considered system is $|-, -, 1\rangle$, the time dependent wave function of the whole system takes the form

$$|\Psi(t)\rangle = Z_{14}|-, -, 1\rangle + Z_{24}|+, -, 0\rangle + Z_{34}|-, +, 0\rangle, \quad (10)$$

where

$$Z_{13} = \frac{1}{\Omega}(\alpha^2 + \cos \Omega t), \quad Z_{23} = -\frac{i}{\Omega} \sin \Omega t, \quad Z_{33} = -\frac{\alpha}{\Omega^2}(1 - \cos \Omega t).$$

At last, if the initial state is $|-, -, 0\rangle$, the time dependent wave function will evolve to

$$|\Psi(t)\rangle = e^{-i\omega t}|-, -, 0\rangle. \quad (11)$$

Now we go back to the theme of this paper. If the initial state for two atoms is (2), then using the equations (4)-(11) one can obtain the density operator for the whole system. Taking a partial trace over the heat bath variable one can obtain that the reduced atomic density operator in the two-atom basis evolves to

$$\rho_A(t) = \begin{pmatrix} \rho_{11} & \rho_{12} & \rho_{13} & \rho_{14} \\ \rho_{12}^* & \rho_{22} & \rho_{23} & \rho_{24} \\ \rho_{13}^* & \rho_{23}^* & \rho_{33} & \rho_{34} \\ \rho_{14}^* & \rho_{24}^* & \rho_{34}^* & \rho_{44} \end{pmatrix}. \quad (11)$$

The matrix elements of (11) are given in Appendix.

For two-qubit system described by the density operator $\rho_A(t)$, a measure of entanglement or negativity can be defined in terms of the negative eigenvalues μ_i^- of partial transpose of the reduced atomic density matrix $\rho_A^{T_1}$ [26],[27]. The partial transpose of the reduced atomic density matrix (11) is

$$\rho_A^{T_1}(t) = \begin{pmatrix} \rho_{11} & \rho_{12} & \rho_{13}^* & \rho_{23}^* \\ \rho_{12}^* & \rho_{22} & \rho_{14}^* & \rho_{24}^* \\ \rho_{13} & \rho_{14} & \rho_{33} & \rho_{34} \\ \rho_{23} & \rho_{24} & \rho_{34}^* & \rho_{44} \end{pmatrix}.$$

The negativity is

$$\varepsilon = -2 \sum \mu_i^-. \quad (12)$$

When $\varepsilon = 0$ two qubits are separable and $\varepsilon > 0$ means the atom-atom entanglement. The case $\varepsilon = 1$ indicates maximum entanglement.

The results of calculations of entanglement parameter (12) for initial atomic states (2) are shown in Fig. 1-3. For all figures we put $\varphi_1 = \varphi_2 = 0$.

3. Results

In Fig. 1 we plot the negativity as a function of gt for coherent and three incoherent atomic states described by formula (2). The curves corresponds to a fixed value of dipole strength $\alpha = 0.5$ and weak thermal two-mode field with $\bar{n} = 0.01$. From these figure we can find that, firstly, entanglement can be induced by thermal field for all initial atomic state. The more significant result is that the atomic entanglement may be induced by thermal field when both the atoms are prepared in their excited states. By contrast, the atomic entanglement does not induced by thermal field when both atoms are trapped in cavity and prepared in their excited states [5]. Secondly, the maximum degree of entanglement is slightly enhanced owing to the atomic coherence comparing the curves in Fig.1(c) and Fig.1(d). With increasing of the mean photon number (see Figs.2(a) and Fg2(b)) the value of atom-atom negativity decreases. But for coherent states this decreasing is much sharper. Such entanglement behavior essentially differs from that for model when both atoms are trapped in cavity and interact with a thermal field. In the last case [16] the entanglement is greatly enhanced due to the initial atomic coherence for thermal cavity with large mean photon numbers. Fig. 3 displays how the evolution of entanglement depends on the dipole-dipole strength for $\bar{n} = 1$. From these figures, we find that entanglement revealed by the solid lines ($\alpha = 0.1$) is stronger than that by the dotted lines ($\alpha = 1$) at most time. The value of atomatom negativity is larger when the dipole interaction between the atoms increases. This dependence is much sharp for incoherent initial atomic state.

4. Conclusion

In this paper we have investigated the effect of the atomic coherence on the entanglement of two dipole coupled two-level atoms when only one atom is trapped in a lossless cavity and interacts with one-mode thermal field, and the other one can be spatially moved freely outside the cavity. We have shown that thermal field can produce atom-atom entanglement for all pure initial atomic

states. The results also show that the atom-atom entanglement can be controlled by changing the system parameters, such as the amplitudes of the polarized atoms, the mean photon numbers of thermal field, and the strength of the dipole interaction (or distance between atoms). We have derived that for considered model the dipole-dipole interaction can produce the appreciable amount of entanglement only for small thermal noise if the atoms are prepared in the coherent states.

Acknowledgments

Work is carried out with financial support of the Ministry of Education and Science of the Russian Federation (the state assignment N 1394).

Appendix: Expressions for density matrix Elements in Eq. (11)

The evident expressions for matrix elements in (11) are

$$\begin{aligned}
\rho_{11} &= |a|^2 \sum_{n=0} p_n |Z_{41,n}|^2 + |b|^2 \sum_{n=1} p_n |Z_{42,n-1}|^2 + bc^* \sum_{n=1} p_n Z_{42,n-1} Z_{43,n-1}^* + \\
&+ cb^* \sum_{n=1} \sum_{n=1} p_n Z_{43,n-1} Z_{42,n-1}^* + |c|^2 \sum_{n=1} p_n |Z_{43,n-1}|^2 + |d|^2 \sum_{n=2} p_n |Z_{44,n-2}|^2, \\
\rho_{12} &= ab^* \sum_{n=1} p_n Z_{41,n} Z_{22,n-1}^* + ac^* \sum_{n=1} p_n Z_{41,n} Z_{23,n-1} + \\
&+ bd^* \sum_{n=2} p_n Z_{42,n-1} Z_{24,n-2} + cd^* \sum_{n=2} p_n Z_{43,n-1} Z_{24,n-2}^* + \\
&+ p_1 (bd^* Z_{42,0} G_{24}^* + cd^* Z_{43,0} G_{24}^*) + p_0 (ab^* Z_{41,0} G_{22}^* + ac^* Z_{41,0} G_{23}^*), \\
\rho_{13} &= ab^* \sum_{n=1} p_n Z_{41,n} Z_{32,n-1} + ac^* \sum_{n=1} p_n Z_{41,n} Z_{33,n-1}^* + \\
&+ bd^* \sum_{n=2} p_n Z_{42,n-1} Z_{34,n-2}^* + cd^* \sum_{n=2} p_n Z_{43,n-1} Z_{34,n-2}^* + \\
&+ p_1 (bd^* Z_{42,0} G_{34}^* + cd^* Z_{43,0} G_{34}^*) + p_0 (ab^* Z_{41,0} G_{32}^* + ac^* Z_{41,0} G_{33}^*), \\
\rho_{14} &= ad^* \sum_{n=2} p_n Z_{41,n} Z_{14,n-2}^* + p_1 ad^* Z_{41,1} G_{14}^* + p_0 ad^* Z_{41,0},
\end{aligned}$$

$$\begin{aligned}
\rho_{22} &= |a|^2 \sum_{n=0} p_n |Z_{21,n}|^2 + |d|^2 \sum_{n=2} p_n |Z_{24,n-2}|^2 + \\
&+ bc^* \sum_{n=1} p_n Z_{22,n-1} Z_{23,n-1}^* + cb^* \sum_{n=1} p_n Z_{23,n-1} Z_{22,n-1}^* + \\
&+ |c|^2 \sum_{n=1} p_n |Z_{23,n-1}|^2 + |b|^2 \sum_{n=1} p_n |Z_{22,n-1}|^2 + \\
&+ p_1 |d|^2 |G_{24}|^2 + p_0 (|b|^2 |G_{22}|^2 + bc^* G_{22} G_{23}^* + p_0 (cb^* G_{23} G_{22}^* |c|^2 |G_{23}|^2), \\
\rho_{23} &= |a|^2 \sum_{n=0} p_n Z_{21,n} Z_{31,n}^* + |d|^2 \sum_{n=2} p_n Z_{24,n-2} Z_{34,n-2}^* + \\
&+ bc^* \sum_{n=1} p_n Z_{22,n-1} Z_{33,n-1}^* + cb^* \sum_{n=1} p_n Z_{23,n-1} Z_{32,n-1}^* + \\
&+ |c|^2 \sum_{n=1} p_n Z_{23,n-1} Z_{33,n-1}^* + |b|^2 \sum_{n=1} p_n Z_{22,n-1} Z_{32,n-1}^* + \\
&+ p[1] |d|^2 G_{24} G_{23}^* + p[0] (|b|^2 G_{22} G_{32}^* + bc^* G_{22} G_{33}^* + cb^* G_{23} G_{32}^* + |c|^2 G_{23} G_{33}^*), \\
\rho_{33} &= |a|^2 \sum_{n=0} p_n |Z_{31,n}|^2 + |b|^2 \sum_{n=1} p_n |Z_{32,n-1}|^2 + bc^* \sum_{n=1} p_n Z_{32,n-1} Z_{33,n-1}^* + \\
&+ cb^* \sum_{n=1} p_n Z_{33,n-1} Z_{32,n-1}^* + |c|^2 \sum_{n=1} p_n |Z_{33,n-1}|^2 + |d|^2 \sum_{n=2} p_n |Z_{34,n-2}|^2 + \\
&+ p_1 |d|^2 |G_{34}|^2 + p_0 (|b|^2 |G_{32}|^2 + bc^* G_{32} G_{33}^* + cb^* G_{33} G_{32}^* + |c|^2 |G_{33}|^2, \\
\rho_{34} &= ab^* \sum_{n=1} p_n Z_{31,n} Z_{12,n-1} + ac^* \sum_{n=1} p_n Z_{31,n} Z_{13,n-1} + \\
&+ bd^* \sum_{n=2} p_n Z_{32,n-1} Z_{14,n-2}^* + cd^* \sum_{n=2} p_n Z_{33,n-1} Z_{14,n-2}^* + \\
&+ p_1 (bd^* Z_{32,0} G_{14}^* + cd^* Z_{33,0} G_{14}^*) + p_0 (ab^* Z_{31,0} G_{12}^* + ac^* Z_{31,0} G_{13}^*) + \\
&+ p_0 (bd^* G_{32} + cd^* G_{33}, \\
\rho_{44} &= |a|^2 \sum_{n=0} p_n |Z_{11}|^2 + |b|^2 \sum_{n=1} p_n |Z_{12,n-1}|^2 + bc^* \sum_{n=1} p_n Z_{12,n-1} Z_{13,n-1}^* + \\
&+ cb^* \sum_{n=1} p_n Z_{13,n-1} Z_{12,n-1}^* + |c|^2 \sum_{n=1} p_n |Z_{13,n-1}|^2 + |d|^2 \sum_{n=2} p_n |Z_{14,n-2}|^2 + \\
&+ p_1 |d|^2 |G_{14}|^2 + p_0 (|b|^2 |G_{12}|^2 + bc^* G_{12} G_{13}^* + cb^* G_{13} G_{12}^* + |c|^2 |G_{13}|^2 + |d|^2),
\end{aligned}$$

where

$$a = \cos \theta_1 \cos \theta_2, \quad b = \cos \theta_1 \sin \theta_2 e^{i\varphi_2}, \quad c = \cos \theta_2 \sin \theta_1 e^{i\varphi_1}, \quad d = \sin \theta_1 \sin \theta_2 e^{i(\varphi_1 + \varphi_2)}.$$

References

References

- [1] M.A.Nielsen M.A., I.L. Chuang, Quantum Computation and Quantum Information, Cambrige: Cambrige University Press, 2000.
- [2] I. Buluta, S. Ashhab, F. Nori, Reports on Progress in Physics 74 (2011) 104401.
- [3] M.B. Plenio, S.F. Huelda, A. Beige, P.L. Knight, Phys. Rev. A. 59 (1999) 2468–2475.
- [4] S. Bose S., I. Fuentes-Guridi, P.L. Knight, V. Vedral, Phys.Rev.Lett. 87 (2001) 050401.
- [5] M.S. Kim, J. Lee, D. Ahn, P.L. Knight, Phys. Rev. A. 65 (2002) 040101.
- [6] Zhang B., Optics Commun. 283 (2010) 4676-4679.
- [7] L. Zhou, H.S. Song, J. Optics B: Quantum Semiclass. Opt. 6 (2004) 378–384.
- [8] L. Zhou, H.S. Song, J. Optics B: Quantum Semiclass. Opt. 4 (2002) 425–429.
- [9] E.K. Bashkistrov, Laser Phys. Lett. 3 (2006) 145–150.
- [10] L.S. Aguiar, P.P. Munhoz, A. Vidiella-Barranco, J.A. Roversi, J. Opt. B7 (2005) S769–771.
- [11] X.-P. Liao, M.-F. Fang, J.-W. Cai, X.-J. Zheng, Chin. Phys. B17 (2008) 2137–2142.
- [12] E.K. Bashkistrov, M.P. Stupatskaya, Laser Phys. 19 (2009) 525–530.
- [13] E.K. Bashkistrov, M.S. Mastyugin, Optics and Spectroscopy 116 (2014) 630–634.

- [14] X.-Y. Wang, X.-S. Chen, J. Phys. B39 (2006) 3805–3814.
- [15] Y.-H. Hu, M.-F. Fang, Q. Wu, Chin. Phys. 16 (2007) 2407–2414.
- [16] Y.-H. Hu, M.-F. Fang, Comm. Theor. Phys. 54 (2010) 421–426.
- [17] Y.-H. Hu, M.-F. Fang, C.-L. Jiang, K. Zeng, Chin. Phys. 17 (2008) 1784–1790.
- [18] F.A.A. El-Orany, J.Opt.Soc.Am. B28 (2011) 2087–2097.
- [19] Y.Q. Guo, H.J. Cao, H.S. Song, arXiv:quant-ph/0509142 (2005) 7 p.
- [20] C. Marr, A. Beige, G. Rempe, Phys.Rev. A68 (2003) 033817.
- [21] S. Mancini, S. Bose, G. Rempe, Phys.Rev. A70 (2004) 022307.
- [22] G. Chimczak, Phys.Rev. A71 (2005) 052305.
- [23] L.T. Shen, X.Y. Chen, Z.B. Yang, H.Z. Wu, S.B. Zheng, Phys.Rev. A84 (2011) 064302.
- [24] E.K. Bashkirov, Y.A. Nikiforova, Computer Optics 36 (2012) 468–473 (In Russian).
- [25] M. Lu, Y. Xia, L.T. Shen, J. Song, N.B. An, Phys.Rev. A89 (2014) 012326.
- [26] A. Peres, Phys. Rev. Lett. 77 (1996) 1413–1415.
- [27] R. Horodecki, M. Horodecki, P. Horodecki, Phys. Lett. A223 (1996) 333–339.







Figure 1: The negativity as a function of gt for the model with $\bar{n} = 0.01$ and $\alpha = 0.5$. The initial atomic state: a) $|\Psi(0)\rangle_A = |+, -\rangle$ (a), $|\Psi(0)\rangle_A = |-, -\rangle$ (b), $|\Psi(0)\rangle_A = |+, +\rangle$ (c) and $|\Psi(0)\rangle_1 = (1/\sqrt{2})(|++\rangle + |--\rangle)$, $|\Psi(0)\rangle_2 = (1/\sqrt{2})(|+-\rangle + -+ \rangle)$ (d).




Figure 2: The negativity as a function of gt for the model with $\bar{n} = 1$ (solid) and $\bar{n} = 2$ (dashed). The initial atomic state: a) $|\Psi(0)\rangle_A = |+, -\rangle$ (a) and $|\Psi(0)\rangle_1 = (1/\sqrt{2})(|+\rangle + |-\rangle)$, $|\Psi(0)\rangle_2 = (1/\sqrt{2})(|+\rangle + |-\rangle)$ (b). The dipole strength $\alpha = 0.5$




Figure 3: The negativity as a function of gt for the model with $\alpha = 0.1$ (solid) and $\alpha = 1$ (dashed). The initial atomic state: a) $|\Psi(0)\rangle_A = |+, -\rangle$ (a) and $|\Psi(0)\rangle_1 = (1/\sqrt{2})(|+\rangle + |-\rangle)$, $|\Psi(0)\rangle_2 = (1/\sqrt{2})(|+\rangle + |-\rangle)$ (b). The mean photon number $\bar{n} = 0.5$